

Evolution of massive black hole binaries in gaseous environments



Universität
Zürich^{UZH}



Alessia Franchini


November 5th 2024

@ LISA AstroWG meeting, Munich (D)

Massive black hole binaries path to coalescence



Credit: NASA/ESA/Hubble




Credit: E. Bortolas

Massive black hole binaries in gaseous environments




After a successful dynamical friction driven inspiral, a bound binary in the central parsec of the remnant host galaxy evolves through the interaction with gas before diving into the GW dominated phase

Pulsar Timing Array (PTA) binaries may reside in massive discs whose self-gravity cannot be neglected

LISA binaries evolve within gaseous discs where the gas self-gravity can be neglected



PTA Massive black hole binaries in gaseous environments







We find binaries embedded in massive self-gravitating discs to always in-spiral while reaching an orbital eccentricity which depends on the initial value and on the initial mass ratio.



LISA Massive black hole binaries in gaseous environments

We use hyper-Lagrangian refinement to investigate the effect of the disc temperature during the interaction between an equal mass circular binary and an isothermal circumbinary disc.





We find binary out-spiral for intermediate values of the disc temperature




LISA Massive black hole binaries electromagnetic counterparts

We use hyper-Lagrangian refinement coupled with 2.5PN corrections to the binary orbit to investigate electromagnetic counterparts of LISA binaries.




LISA Massive black hole binaries multi band light curves

We use hyper-Lagrangian refinement coupled with 2.5PN corrections to the binary orbit to investigate electromagnetic counterparts of LISA binaries prior, during and post merger.



$H/R = 0.1$




$H/R = 0.03$




Post merger signatures of LISA massive black hole binaries

We investigated the accretion rate onto the merger remnant in two kick scenarios.



Remnant moves along the z-axis, dragging the disc outside its initial orbital plane


Remnant moves along the y-axis, towards the cavity pericentre



$H/R = 0.03$





Chirping of massive black hole binaries in the optical band



Spectrogram of in-spiral electromagnetic signal in the optical band for circular equal mass binaries.

Deviation from the GW driven inspiral rate

Modification to the orbital evolution due to the presence of gas around the binary.



- Binary inspiral on a shorter timescale (~ 2 hours)
- Phase shift ~ 0.14 radians

Conclusions

- Binary semi-major axis decreases with time as a result of its interaction with a circumbinary disc in a vast region of the parameter space
- PN corrections to the binary dynamics allow us to extract electromagnetic signatures prior, during and after the merger
- We find significant orbital phase deviation with respect to the evolution of the binary in vacuum (see also **Mudit Garg's talk**)
- Merger characterised by a ~ 2 orders of magnitude decrease in the X-ray flux followed by an increase in the UV flux
- Off plane kicks are characterised by a very fast increase of the accretion rate post merger
- Periodicity on the binary orbital period and lump modulation strength depends on the binary parameters but it is difficult to detect with a handful of binary orbits (**see Fabiola Cocchiara's talk**)
- The chirping signal can in principle be detected also in the optical band